

Amendment dated December 22, 2005
Atty Dkt No. 1950-0001
Client No. CIT-3113
Serial No.: 10/091,372

Exhibit A

[Sign in](#)



[Web](#) [Images](#) [Groups](#) [News](#) [Froogle](#) [Local](#)^{New!} [more »](#)

"tertiary nitrogen atoms"

[Search](#)

[Advanced Search](#)
[Preferences](#)

Web

Results 1 - 10 of about 178 for "**tertiary nitrogen atoms**". (0.34 seconds)

[\[PDF\] Characterization of Co\(II\) Binding to PAMAM G4-NH Dendrimer in ...](#)

File Format: PDF/Adobe Acrobat - [View as HTML](#)

in the terminal branches, and there are **tertiary nitrogen atoms** that are ...

approximately 4 **tertiary nitrogen atoms** involved for every Co(II) bound to the ...

[www.cee.org/rsi/docs/2004Papers/al_rabe-khaled-caltech-oral.pdf](#) - [Similar pages](#)

[\[PDF\] Influence of Process Factors on the Structure of Urea Formaldehyde ...](#)

File Format: PDF/Adobe Acrobat

tertiary nitrogen atoms (k). Signals belonging to the middle groups in the

polymethylene ... groups at **tertiary nitrogen atoms** (i) and di- and trisub- ...

[www.ingentaconnect.com/content/maik/rjac/2003/00000076/00000003/00475229](#) - [Similar pages](#)

[Chemistry Journals published by the Korean Chemical Society and ...](#)

Effect of Uncoordinated **Tertiary Nitrogen Atoms** in Hexaaza Macrocyclic Nickel(II)

Complexes on Axial Binding of Anions and Water in Acidic Solutions ...

[newjournal.kcsnet.or.kr/.../j_bkcs_archives_](#)

[view2.htm?code=B950308&qpage=j_bkcs&spage=j_bkcs_01](#) - 33k - [Cached](#) - [Similar pages](#)

[Tertiary amines having carbonate and urethane groups - Patent 5834579](#)

... of polyurethane foams is preferably selected so that from about 0.01 to about

1% by weight **tertiary nitrogen atoms** are present in the reaction mixture. ...

[www.freepatentsonline.com/5834579.html](#) - 46k - [Cached](#) - [Similar pages](#)

[Full Text Button for patent number 6673890](#)

... comprising: a linear or branched hydrophobic polyamine backbone having 2 to 10 tertiary amino nitrogen atoms and a spacer between two **tertiary nitrogen atoms**, ...

[www.uspto.gov/web/patents/patog/week01/OG/html/1278-1/US06673890-20040106.html](#) - 12k -

[Supplemental Result](#) - [Cached](#) - [Similar pages](#)

[\[PDF\] molecules](#)

File Format: PDF/Adobe Acrobat - [View as HTML](#)

Six imidazole molecules are coordinated through their **tertiary nitrogen atoms**

to each nickel (II) ion, and two dtp anions are out of ...

[www.mdpi.org/molecules/papers/81200866.pdf](#) - [Similar pages](#)

[Amersham Biosciences - Receptor Internalization](#)

... Unlike classic cyanine dyes, which contain **tertiary nitrogen atoms** in both the heterocyclic rings, the pH-sensitive dyes contain secondary nitrogen atoms in ...

[www.amershambiosciences.com.cn/aptrix/upp00919.nsf/](#)

[Content/drugscr_applications%5Cdrugscr_applications_gp...](#) - 31k - [Supplemental Result](#) -

[Cached](#) - [Similar pages](#)

[Anhydrous Proton-Conducting Membranes for Fuel Cells](#)

This work has included an investigation of acid salts of triethylenediamine in

which each molecule contains two **tertiary nitrogen atoms** that can be ...

[www.nasatech.com/Briefs/aug05/NPO30493.html](#) - 8k - [Cached](#) - [Similar pages](#)

[PDF] [Synthesis and characterization of polypyrrolepolyurethane ...](#)

File Format: PDF/Adobe Acrobat

cursor with **tertiary nitrogen atoms** in the polymer chain was prepared ... tion of **tertiary nitrogen atoms** to acrylic acid, with appearance of ion pairs ...
doi.wiley.com/10.1002/apmc.1997.052450112 - [Similar pages](#)

[Biodegradability of ethylenediamine-based complexing agents and ...](#)

Tetra(penta)substituted derivatives with two or more **tertiary nitrogen atoms** and carboxymethyl or 2-hydroxyethyl groups in the molecule (EDTA, DTPA, PDTA, ...
www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&
db=PubMed&list_uids=11482674&dopt=Abstract - [Similar pages](#)

Try searching for "**tertiary nitrogen atoms**" on [Google Book Search](#)

Goooooooooooooogle ▶

Result Page: 1 2 3 4 5 6 7 8 9 [Next](#)



Free! Instantly find your email, files, media and web history. [Download now.](#)

"tertiary nitrogen atoms"

[Search](#)

[Search within results](#) | [Language Tools](#) | [Search Tips](#) | [Dissatisfied? Help us improve](#)

[Google Home](#) - [Advertising Programs](#) - [Business Solutions](#) - [About Google](#)

©2005 Google

USPTO PATENT FULL-TEXT AND IMAGE DATABASE[Home](#)[Quick](#)[Advanced](#)[Pat Num](#)[Help](#)[Next List](#)[Bottom](#)[View Cart](#)

Searching 1976 to present...
































Results of Search in 1976 to present db for:





"tertiary nitrogen atoms": 435 patents.


Hits 1 through 50 out of 435

[Next 50 Hits](#)[Jump To](#)[Refine Search](#)

PAT. NO.	Title
1 6,964,986	Aqueous adhesive system, method for the production and use thereof
2 6,956,086	Water dispersible epoxy resins
3 6,916,954	Catalyst system and method for carbonylation
4 6,916,767	Antioxidant amines based on n-(4aniliophenyl) amides antioxidant amines based on n-(4-anilinophenyl) amides
5 6,911,558	Method for purifying chloromethyl chloroformate
6 6,897,266	Dendritically modified polyurethanes
7 6,875,790	Cyclic sulfone containing retroviral protease inhibitors
8 6,875,245	Ester-terminated poly(ester-amides) in personal care products
9 6,864,349	Aqueous suspensions containing polymerized fatty acid-based polyamides
10 6,849,709	Aerothricin derivatives
11 6,786,940	Paraffin dispersants with a lubricity effect for distillates of petroleum products
12 6,777,530	Use of crosslinked nitrogenous compounds which are soluble or dispersible in water in detergents and cleaners
13 6,765,035	Process for making rigid and flexible polyurethane foams containing a fire-retardant
14 6,762,260	Organoborane amine complex polymerization initiators and polymerizable compositions
15 6,746,542	Modifying starch with cationic polymers and use of the modified starches as dry-strength agent

- 16 [6,740,716](#)  [Organoborane amine complex polymerization initiators and polymerizable compositions](#)
- 17 [6,740,633](#)  [Polyelectrolyte complexes and a method for production thereof](#)
- 18 [6,723,882](#)  [Preparation of dialkyl ketones](#)
- 19 [6,703,523](#)  [Zwitterionic polyetherpolyamines and process for their production](#)
- 20 [6,673,890](#)  [Zwitterionic polyamines and process for their production](#)
- 21 [6,653,436](#)  [Water dispersible epoxy resins](#)
- 22 [6,649,029](#)  [Nonfelting wool and antifelt finishing process](#)
- 23 [6,641,655](#)  [Fluidizing agents](#)
- 24 [6,616,797](#)  [Preparation of adhesive bonds using carbodiimide-containing dispersions with heat activation](#)
- 25 [6,596,678](#)  [Laundry detergent compositions containing a polymer for fabric appearance improvement](#)
- 26 [6,592,857](#)  [Tertiary amide terminated polyamides in cosmetics](#)
- 27 [6,569,361](#)  [Electrochrome polymer systems](#)
- 28 [6,552,160](#)  [Ester-terminated poly\(ester-amides\) useful for formulating transparent gels in low polarity fluids](#)
- 29 [6,551,761](#)  [Photoactivatable nitrogen-containing bases based on .alpha.-ammonium ketones, iminium ketones or amidinium ketones and aryl borates](#)
- 30 [6,548,461](#)  [Polyalkene alcohol polyetheramines and their use in fuels and lubricants](#)
- 31 [6,503,522](#)  [Tertiary amide-terminated polyamides in structured personal care compositions](#)
- 32 [6,503,077](#)  [Gelled articles containing tertiary amide-terminated polyamide](#)
- 33 [6,489,440](#)  [Cyclic compounds](#)
- 34 [6,489,374](#)  [Photoactivatable bases containing nitrogen](#)
- 35 [6,469,131](#)  [Structured composition containing tertiary amide-terminated polyamide for personal care products](#)
- 36 [6,468,652](#)  [Elastic polyurethane-urea fiber and method for producing the same](#)
- 37 [6,359,037](#)  [Polyamine/epoxy-functional amidoamine product with epoxy resin](#)
- 38 [6,346,204](#)  [Process for preparing rigid and flexible polyurethane foams](#)
- 39 [6,333,134](#)  [Multilayered photopolymer element including sensitivity controlling agents](#)
- 40 [6,313,257](#)  [Poly \(mercaptopropylaryl\) curatives](#)
- 41 [6,277,928](#)  [Epoxy-functional amidoamine reacted with excess polyamine and monoepoxy as epoxy curative](#)
- 42 [6,268,466](#)  [Tertiary amide terminated polyamides and uses thereof](#)
- 43 [6,265,118](#)  [Image element and image formation method](#)
- 44 [6,242,509](#)  [Gels including bioactive components](#)
- 45 [6,235,931](#)  [Partial oxidation of polyoxyalkylene polyol compositions to polycarboxylic acid compositions](#)
- 46 [6,228,828](#)  [Laundry detergent compositions with anionically modified, cyclic amine based polymers](#)

- 47 [6,221,954](#)  [Cationic polyurethane compositions, quaternary ammonium salts and methods for their preparation](#)
- 48 [6,221,934](#)  [Aqueous dispersions of epoxy resins](#)
- 49 [6,217,687](#)  [Processes for treating a surface of a thermoplastic resin film](#)
- 50 [6,211,258](#)  [Process for preparing rigid and flexible polyurethane foams](#)
-



Synthesis and Crystal Structure of Hexakis(imidazole) nickel (II) O,O'-diphenyldithiophosphate [Ni(Im)₆](Ph₂O₂PS₂)₂

Shusheng Zhang *, Shiying Wang, Yonghong Wen and Kui Jiao

College of Chemistry and Molecule Engineering, Qingdao University of Science and Technology, Qingdao, Shandong 266042, P. R. China.

* Author to whom correspondence should be addressed. E-mail: zhangshush@public.qd.sd.cn; Tel: +86-532-4022750; Fax: +86-532-4023927.

Received: 10 October 2003; in revised form: 24 November 2003 / Accepted: 28 November 2003 / Published: 31 December 2003.

Abstract: The crystal and molecular structures of [Ni(Im)₆](dtp)₂ (Im = imidazole, dtp = O,O'-diphenyldithiophosphate) have been determined by X-ray crystallography. It crystallizes in the triclinic system, space group P $\bar{1}$, with cell parameters $a = 9.375(2)$, $b = 12.324(3)$, $c = 13.285(3)$ Å, $\alpha = 107.86(3)$, $\beta = 102.28(3)$, $\gamma = 109.24(3)$, and $Z = 1$. The crystal structure of the title compound is built up of discrete monomeric molecules of [Ni(Im)₆](dtp)₂. The nickel (II) ion is hexacoordinated by six imidazole molecules and the coordination environment of Ni (II) is of octahedral geometry. In the solid state, a network of N-H...S intermolecular hydrogen bonds connect the Ni(Im)₆ moieties and O,O'-diphenyldithiophosphate molecules, forming a three-dimensional structure.

Keywords: Imidazole complexes; O,O'-diphenyl dithiophosphate; crystal structure; hydrogen bonds; three-dimensional network

Introduction

Imidazole is of considerable interest as a ligand in that its presence in many biological systems provides a potential binding site for metal ions [1, 2]. Imidazole itself is usually a unidentate ligand and forms complexes with metal ions through its tertiary nitrogen atom. Some complexes of imidazole and its derivatives with transition-metal ions have been reported [3-5].


Complexes of nickel (II) and imidazole ligand have been studied as models for copper proteins that contain both functionalities in the side chain [6]. Some of these nickel (II) complexes were found to exhibit a variety of pharmacological activity and superoxide dismutase activities [7]. The adducts and their formation reactions have also been found useful in a variety of ways, such as in biological applications and as both ultraviolet absorbers and antioxidants. Recently, the structure of hexakis(imidazole) nickel (II) disalicylate, $[\text{Ni}(\text{Im})_6](\text{sal})_2$, has been reported [8]. In this paper, we describe the synthesis, and crystal structure of the title compound, hexakis(imidazole) nickel (II) O,O'-diphenyldithiophosphate, $[\text{Ni}(\text{Im})_6](\text{Ph}_2\text{O}_2\text{PS}_2)_2$.

Results and Discussion

X-ray Crystal Structure of the Title Compound

The X-ray structure of the complex $[\text{Ni}(\text{Im})_6](\text{dtp})_2$ is built up of discrete monomeric molecules. Figure 1 shows a perspective view of the title compound with atomic numbering scheme and Figure 2 a perspective view of the crystal packing in the unit cell. Table 1 contains atomic positions and equivalent temperature factors for non-hydrogen atoms. Selected bond lengths and angles are presented in Table 2.


Figure 1 Molecular structure for $[\text{Ni}(\text{Im})_6](\text{dtp})_2$ with the atomic numbering scheme



The crystal structure of the complex consists of $[\text{Ni}(\text{Im})_6]^{2+}$ cation and two dtp anions, linked by electrostatic forces and hydrogen bonds (shown as dashed lines in Figure 2). The coordination model of the nickel (II) can be described as being of octahedral geometry. Six imidazole molecules are coordinated through their tertiary nitrogen atoms to each nickel (II) ion, and two dtp anions are out of

the coordination sphere and balance the charge. The bond distances of Ni(1)-N(1), Ni(1)-N(3) and Ni(1)-N(5) of $[\text{Ni}(\text{Im})_6](\text{dtp})_2$ are 2.133(3), 2.132(3) and 2.125(3) Å, respectively. These distances are in agreement with those reported for octahedral $[\text{Ni}(\text{Im})_6]^{2+}$ complex [2.120(1), 2.124(1), 2.141(1) Å], $[\text{Ni}(\text{Im})_6](\text{sal})_2$ (sal = salicylate) [8]. In the solid state, one dtp sulfur atom forms a N-H \cdots S intermolecular hydrogen bond with the nitrogen atom of $\text{Ni}(\text{Im})_6^{2+}$ (Figure 2), and the donor and acceptor distances are N(2) \cdots S(1) 3.6547, C(17) \cdots S(2) (2-x, 1-y, 1-z) 3.3052, C(11) \cdots S(2) (2-x, 1-y, 1-z) 3.7945 and C(20) \cdots S(1) (2-x, -y, 1-z) 3.3807 Å (Table 3), the whole complexes forming a 3-dimensional network structure.

Figure 2 A view of the crystal packing down the z axis for $[\text{Ni}(\text{Im})_6](\text{dtp})_2$. The dashed lines represent hydrogen bonds between dtp anions and imidazoles.



IR Spectroscopy

The IR spectrum of the complex shows characteristic absorption bands at 3401 and 552 cm^{-1} , assigned to $\nu(\text{N-H})$ and $\nu(\text{Ni-N})$ vibrations, respectively. This indicates that the atom coordinated with Ni(II) is the double bond nitrogen ($=\text{N}$) in the imidazole molecule. This result is in agreement with that of X-ray crystal structure determination.

Table 1 Atomic coordinates and equivalent isotropic displacement parameters

Atom	x	y	z	U_{eq}
Ni1	0.50000	0.00000	0.50000	0.03408
S1	-0.02504	0.11075	0.11362	0.05583
S2	0.23540	0.33035	0.36706	0.04651
P1	0.11983	0.28764	0.20898	0.04016
O1	0.25663	0.34717	0.16152	0.05387
O2	0.02372	0.37457	0.19719	0.05210
N1	0.70819	0.07677	0.45909	0.04516
N2	0.86587	0.12438	0.36401	0.07807
N3	0.49052	0.17820	0.55780	0.04184
N4	0.44719	0.34158	0.64827	0.09284
C1	0.22540	0.34804	0.05361	0.06202
C2	0.20081	0.44556	0.03942	0.09705
C3	0.17904	0.45204	-0.06446	0.14530
C4	0.18613	0.36170	-0.15079	0.15496
C5	0.20809	0.25992	-0.13735	0.13347
C6	0.22857	0.25275	-0.03206	0.09396
C7	-0.10835	0.36540	0.23377	0.05025
C8	-0.25918	0.27493	0.16484	0.07322

U_{eq} is defined as one third of the trace of the orthogonalized U_{eq} tensor.

Table 2 Selected bond distances (Å) and bond angles (°) of title compound

Ni1-N1	2.133(3)	S1-P1	1.955 (2)
Ni1-N3	2.132(3)	S2-P1	1.959 (2)
Ni1-N5	2.125(3)	P1-O1	1.609(2)
C1-C2	1.355(7)	P1-O2	1.632(2)
C1-C6	1.376(7)	O1-C1	1.405(4)
O1-P1-O2	97.47(13)	C1-C2-C3	119.7(6)
O1-P1-S2	105.29(11)	C4-C3-C2	119.6(7)
S1-P1-S2	117.31(7)	N5-Ni1-N3	90.40(11)
N5-Ni1-N1	90.81(11)	N3-Ni1-N1	91.05(11)
C14-C13-N1	109.2(4)	N1-C15-N2	111.6(4)

Symmetry transformations used to generate equivalent atoms: \$1 -x+1, -y, -z+1

Table 3 Hydrogen bond distances (Å) and angles of title compound

D	H	A	symm	D-H	H...A	D...A	D-H...A Bond angle
N(2)	H(8A) (2a)	S(1)		0.9287	2.7403	3.6547	168.35
C(17)	H(4A)	S(2)	2766	0.9224	2.3836	3.3052	177.03
C(11)	H(11A)	S(2)	2766	0.9467	2.8717	3.7945	165.15
C(20)	H(20A)	S(1)	2756	0.9170	2.5156	3.3807	157.42

[2766]=2-x,1-y,1-z; [2756]=2-x,-y,1-z

Acknowledgements

We thank the National Natural Science Foundation of China (Project 20275020) and the Natural Science Foundation of Shandong Province (Project Z2002B02) for financial support.

Experimental

General

All commercially available chemicals were of analytical reagent grade and used directly without further purification. The C, N and H contents were determined by using an Elementar Vario EL analyzer. Infrared spectra were recorded from KBr pellets on a Midac Prospekt IR spectrometer. All operations were carried out under ambient conditions.

Synthesis of the complex $[\text{Ni}(\text{Im})_6](\text{dtp})_2$

To a warm solution of excess imidazole (1.0 g, 15 mmol) in EtOH (50 mL) was added with stirring nickel (II) O, O'-diphenyl dithiophosphate, and the mixture was refluxed for 40 min. The blue solution was filtered and the filtrate was left to stand undisturbed. Upon slow evaporation at room temperature, a blue crystalline solid suitable for X-ray structure determination appeared several weeks later and was separated by filtration. Main IR absorption bands (KBr, cm^{-1}): 3401 (N-H), 1491 (C=N), 1591, 1530 (phenyl ring stretch), 552 (Ni-N). Anal. Data: Calcd for $\text{C}_{42}\text{H}_{44}\text{N}_{12}\text{O}_4\text{P}_2\text{S}_4\text{Ni}_1$ C, 48.94; H, 4.27; N, 16.31 %. Found: C, 48.93; H, 4.30; N, 16.21 %.

Crystallographic Data and Structure Determination [9]

The selected crystal of $[\text{Ni}(\text{Im})_6](\text{dtp})_2$ was mounted on a Nonius CAD4 diffractometer, reflection data for the unit cell determination were measured at 293 K using $\text{MoK}\alpha$ radiation ($\lambda = 0.71073 \text{ \AA}$) with a graphite monochromator. The technique used was ω -scan with $0.995 < \theta < 25.04^\circ$ for the complex. Absorption correction was made with π -scans. The structure was solved by direct method

and refined by full-matrix least-squares method on F^2_{obs} by using the SHELXTL software package [10]. All non-H atoms were anisotropically refined. The hydrogen atoms were geometrically fixed and allowed to ride on the parent atoms to which they are attached. The final conventional $R = 0.0604$, $wR = 0.1388$ for 3843 reflections with $I > 2\sigma(I)$; $w = 1/[\sigma^2(F^2_o) + (0.0600P)^2 + 1.4652P]$ where $P = (F^2_o + 2(F^2_c))/3$. The molecular graphics were created by SHELXTL. Atomic scattering factors and anomalous dispersion correction were taken from International Table for X-Ray Crystallography [11]. A summary of the key crystallographic information is given in Table 4.

Table 4 Summary of crystallographic results for $[\text{Ni}(\text{Im})_6](\text{dtp})_2$

Formula	$\text{C}_{42}\text{H}_{44}\text{N}_{12}\text{O}_4\text{P}_2\text{S}_4\text{Ni}_1$
Formula weight	1029.78
Color/shape	Blue/prism
Crystal system	Triclinic
Space group	$P\bar{1}$
$a/\text{\AA}$	9.375(2)
$b/\text{\AA}$	12.324(3)
$c/\text{\AA}$	13.285(3)
$\alpha/^\circ$	107.86(3)
$\beta/^\circ$	102.28(3)
$\gamma/^\circ$	109.24(3)
$V/\text{\AA}^3$	1292(5)
Z value	1
$D_{\text{calcd.}}/\text{g.cm}^{-3}$	1.307
μ/mm^{-1}	0.65
$F(000)$	534
Crystal size/mm	$0.20 \times 0.24 \times 0.20$
Temp./K	293(2)
θ ranges/ $^\circ$	0.995–25.04
$h/k/l$	–10,11/–14,0/–15,15
Reflections collected	4551
Independent reflections	3843
Absorption correction	Empirical
No. restraints/ No. parameters	0/295
GOF	1.085
Final R indices [$I > 2\sigma(I)$]	$R_1=0.0473$ $wR_2=0.1246$
R indices(all data)	$R_1=0.0604$ $wR_2=0.1388$
Largest peak and hole/ $\text{e}^{-}\text{nm}^{-3}$	660 and –490

References and Notes

- 1 Brooks P.; Davidson N. *J. Amer. Chem. Soc.* **1960**, *82*, 2118.
- 2 Davis, W. J.; Smith, J. *J. Chem. Soc. (A)* **1971**, 317.
- 3 Inova M.; Kishita M.; Kabo M. *Bull. Chem. Soc. Jpn.* **1969**, *39*, 1352.
- 4 Eillbeck W.J.; Holmes F.; Underhill A.E. *J. Chem. Soc. (A)* **1967**, 757.
- 5 Xu, L. Z.; Zhao, P. S.; Zhang, S. S. *Chin. J. Chem.* **2001**, *19*, 436.
- 6 Bernarducci E. E.; Bharadwaj. P. K. *Inorg. Chem.* **1983**, *22*, 3911.
- 7 Tamura H.; Imai H.; Kuwahra J.; Sugiura Y. *J. Amer. Chem. Soc.* **1987**, *109*, 6870.
- 8 Jian, F. F.; Wang, Z.X.; Bai, Z. P.; You, X. Z. *J. Chem. Cryst.* **1999**, *29*, 359.
- 9 CCDC 212159 contains the supplementary crystallographic data for this paper. These data can be obtained free of charge via www.ccdc.cam.ac.uk/conts/retrieving.html (or from the CCDC, 12 Union Road, Cambridge CB2 1EZ, UK; fax: +44 1223 336033; e-mail: deposit@ccdc.cam.ac.uk).
- 10 Sheldrick, G. M. *SHELXTL V5 Reference Manual*, Siemens Analytical X-ray Systems, Inc., Madison: Wisconsin (USA), **1996**.
- 11 Wilson, A.J. *International Table for X-Ray Crystallography*, Vol C; Kluwer Academic Publishers: Dordrecht, **1992**.

Samples Availability: Available from the corresponding author.

© 2003 by MDPI (<http://www.mdpi.org>). Reproduction is permitted for noncommercial purposes.